

Supplementary Table 2. Multiple logistic regression in clinically normotensive men and women (combined) without a history of hypertension according to the presence of electrocardiographic left ventricular hypertrophy or normal sinus rhythm

Variable	No. of subjects with information	Adjusted odds ratio (95% CI)
Model A	2,173	
Male sex (ref: female)		3.06 (1.67–5.59)
Age		1.03 (1.00*–1.06)
SBP		1.05 (1.02–1.07)
Obesity [†] (ref: no obesity)		0.54 (0.30–0.97)
Monthly household income (10,000 won/mo)		
<200		1.00
≥200		0.80 (0.30–2.13)
Model B	2,380	
Male sex		1.71 (0.75–3.86)
Age		1.03 (1.00*–1.06)
SBP		1.05 (1.03–1.07)
Obesity		0.59 (0.34–1.01)
Smoking		
Never		1.00
Ex-smoker		2.03 (0.93–4.41)
Current		2.05 (0.94–4.46)
Model C	1,381	
Male sex		4.82 (1.94–11.99)
Age		1.05 (1.01–1.09)
SBP		1.05 (1.01–1.08)
Obesity		0.49 (0.21–1.11)
Regular exercise [‡]		1.74 (0.84–3.60)
Model D	2,582	
Male sex		3.39 (1.91–6.03)
Age		1.03 (1.00*–1.05)
SBP		1.05 (1.02–1.07)
Obesity		0.58 (0.33–1.00*)
History of stroke		2.41 (0.52–11.16)
Model E	2,582	
Male sex		3.38 (1.90–6.02)
Age		1.03 (1.00*–1.05)
SBP		1.05 (1.02–1.07)
Obesity		0.58 (0.34–1.01)
History of dyslipidemia		1.07 (0.48–2.42)
Model F	2,582	
Male sex		3.31 (1.86–5.90)
Age		1.03 (1.00–1.05)
SBP		1.05 (1.02–1.07)
Obesity		0.56 (0.33–0.98)
Pulse rate		0.97 (0.94–1.00)
Model G	2,581	
Male sex		3.45 (1.93–6.15)
Age		1.03 (1.00*–1.06)
SBP		1.05 (1.02–1.07)
Obesity		0.59 (0.34–1.03)
Fasting blood glucose		1.00 (0.98–1.01)
Model H	2,580	
Male sex		3.27 (1.69–6.30)
Age		1.03 (1.00*–1.06)
SBP		1.05 (1.02–1.07)
Obesity		0.58 (0.33–1.01)
Uric acid		1.02 (0.83–1.25)

(Continued to the next page)

Supplementary Table 2. Continued

Variable	No. of subjects with information	Adjusted odds ratio (95% CI)
Model I	2,580	
Male sex		3.53 (1.65–7.56)
Age		1.03 (1.00*–1.05)
SBP		1.05 (1.02–1.07)
Obesity		0.58 (0.34–1.01)
hemoglobin		0.98 (0.78–1.23)
Model J	2,580	
Male sex		3.33 (1.87–5.93)
Age		1.03 (1.00–1.05)
SBP		1.05 (1.02–1.07)
Obesity		0.57 (0.33–0.98)
Aspartate aminotransferase		1.01 (0.99–1.02)

Data describing medical histories and life style were obtained from structured self-questionnaires.

CI, confidence interval; ref, reference; SBP, systolic blood pressure.

*Upper or lower limit of CI=1.00 and P<0.05. [†]Body mass index ≥25 kg/m². [‡]At least 150 min of moderate-intensity aerobic physical activity throughout the week or at least 75 min of vigorous-intensity aerobic physical activity throughout the week.